


MIXING AND INJECTION INSTRUCTIONS FOR PEPTIDES




Remove plastic caps (or peel off metal center piece) from vial of Peptide and Bacteriostatic water.

Wipe down exposed rubber stoppers with a rubbing alcohol pad.

Insert Mixing syringe needle into vial of bacteriostatic water and withdraw suggested amount of solution.

Gently inject bacteriostatic water into vial containing the peptide powder at an angle so that solution slides slowly down the side of the glass into the powder.

Use orange 1cc U-100 insulin syringe to withdraw your suggested dose and inject into abdomen area as specified below.



Uncover the abdomen to see the whole area. Find the waist area. You may give a shot bounded by these landmarks: below the waist, to just above the hip bone, and from where the body curves at the side to about 2 inches from the middle of the abdomen. Avoid the bellybutton. Use the natural line in the middle of the body as a marker.

Make sure you refrigerate the vial.

For more information please contact our concierge team for assistance at
805-608-4582 or reception@amisantabarbara.com

DOSING INSTRUCTIONS FOR PEPTIDES

Epitalon

25mg Vial: Reconstitute w/ 5ml BWFI

Ipamorelin

10mg Vial: Reconstitute w/ 5ml BWFI

Melanotan

10mg Vial: Reconstitute w/ 5ml BWFI

Mots C

40mg Vial: Reconstitute w/ 5ml BWFI

NAD+

100mg Vial: Reconstitute w/ 10ml BWFI

Pentosan

10mg Vial: Reconstitute w/ 5ml BWFI

PT-141

20mg Vial: Reconstitute w/ 2ml BWFI

Tesamorelin

5mg Vial: Reconstitute w/ 5ml BWFI

Tirzepatide

40mg Vial: Reconstitute w/ 4 mL BWFI

Thymosin Alpha

10mg Vial: Reconstitute w/ 5ml BWFI

Thymosin Beta 10

10mg Vial: Reconstitute w/ 5ml BWFI

Semaglutide

7mg Vial: Reconstitute w/ 3.5ml BWFI

Semaglutide w/ BPC

7mg Vial: Reconstitute w/ 4.5ml BWFI

****BWFI = Bacteriostatic Water For Injections***

For more information please contact our concierge team for assistance at
805-608-4582 or reception@amisantabarbara.com